

Right-handed Majorana Neutrino Mass Matrices for Generating Bimaximal Mixings in Degenerate and Inverted Models of Neutrinos

Mahadev Patgiri^{†1} and N. Nimai Singh[‡]

[†]*Department of Physics, Cotton College, Guwahati-781001, India*

[‡]*Department of Physics, Gauhati University, Guwahati-781014, India*

Abstract

An attempt is made to generate the bimaximal mixings of the three species of neutrinos from the textures of the right-handed Majorana neutrino mass matrices. We extend our earlier work in this paper for the generation of the nearly degenerate as well as the inverted hierarchical models of the left-handed Majorana neutrino mass matrices using the non-diagonal textures of the right-handed Majorana neutrino mass matrices and the diagonal form of Dirac neutrino mass matrices, within the framework of the see-saw mechanism in a model independent way. Such Majorana neutrino mass models are important to explain the recently reported result on the neutrinoless double beta decay ($0\nu\beta\beta$) experiment, together with the earlier established data on LMA MSW solar and atmospheric neutrino oscillations.

PACS numbers: 12.10.Dm;12.15.Ff;14.60.Gh

Keywords: Majorana neutrino mass, degenerate, inverted hierarchical, mixing angles, mass eigenvalues.

¹corresponding author.
e-mail address: mahadev@scientist.com

1 Introduction

In the context of the recently reported experimental result on double beta decay[1], together with the earlier experimental data on the atmospheric[2] and solar[3] neutrino oscillations, it is important to construct theoretical models which predict the degenerate and inverted hierarchical patterns of the Majorana neutrino mass matrices within the framework of the grand unified theories(GUTs) with or without supersymmetry[4,5]. In this short paper we attempt to generate the degenerate as well as the inverted hierarchical pattern of the left-handed Majorana neutrino mass matrices using the see-saw formula in a model independent way. This is, in fact, a continuation of our earlier work[6] where the neutrino mixings are provided from the texture of the right-handed Majorana mass matrix M_{RR} , while keeping the Dirac neutrino mass matrix m_{LR} in the diagonal form. We had taken the Dirac neutrino mass matrix m_{LR} as either the charged lepton mass matrix ($m_{LR} = \tan\beta m_l$ referred to as case(i)) or the up-quark mass matrix ($m_{LR} = m_{up}$ referred to as case(ii))[7]. While referring to the earlier paper[6] for details, the model successfully generated both the hierarchical and the inverted hierarchical (having opposite sign mass eigenvalues) neutrino mass matrices as a result of the proper choice of the parameters in texture of M_{RR} . In section 2 we present the generation of the degenerate as well as inverted hierarchical neutrino mass matrices using the see-saw formula, and their predictions on mass eigenvalues and mixing angles. Section 3 is devoted to summary and conclusion.

2 Neutrino mass matrices from see-saw formula

The left-handed Majorana neutrino mass matrix m_{LL} is given by the celebrated see-saw formula[8],

$$m_{LL} = -m_{LR}M_{RR}^{-1}m_{LR}^T \quad (1)$$

where m_{LR} is the Dirac neutrino mass matrix in the left-right (LR) convention[9]. The leptonic (MNS) mixing matrix is now given by $V_{MNS} = V_{\nu L}^\dagger$ where $m_{LL}^{diag} = V_{\nu L} m_{LL} V_{\nu L}^T$. Here both m_{LR} and the charged lepton mass matrix m_l are taken as diagonal, whereas the right-handed Majorana neutrino mass matrix M_{RR} as non-diagonal. Using the see-saw formula (1) we

generate both patterns of m_{LL} viz., (I) nearly degenerate and (II) inverted hierarchical neutrino mass models. We concentrate here only on the cases which have bimaximal mixings listed in Table-I.

Table-I: Zeroth order neutrino mass matrices with texture zeros corresponding to the LMA MSW solution with bimaximal mixings [4,10].

<i>Type</i>	m_{LL}	m_{LL}^{diag}
I(A)	$\begin{pmatrix} 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{\sqrt{2}} & -\frac{1}{2} & \frac{1}{2} \end{pmatrix} m_0$	$Diag(1, -1, 1)m_0$
I(B)	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} m_0$	$Diag(1, 1, 1)m_0$
I(C)	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} m_0$	$Diag(1, 1, -1)m_0$
II(A)	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{pmatrix} m_0$	$Diag(1, 1, 0)m_0$
II(B)	$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} m_0$	$Diag(1, -1, 0)m_0$

The Dirac neutrino mass matrix m_{LR} involved in the see-saw formula Eq.(1), can be either the charged lepton mass matrix m_l (case (i)) or the up-quark mass matrix m_{up} (case (ii)) depending on the particular SUSY SO(10) GUT model and the contents of the Higgs fields employed[6,7]:

$$m_{LR} = \tan\beta \begin{pmatrix} \lambda^6 & 0 & 0 \\ 0 & \lambda^2 & 0 \\ 0 & 0 & 1 \end{pmatrix} m_\tau, \quad (2)$$

and

$$m_{LR} = \begin{pmatrix} \lambda^8 & 0 & 0 \\ 0 & \lambda^4 & 0 \\ 0 & 0 & 1 \end{pmatrix} m_t, \quad (3)$$

respectively. The above two forms of m_{LR} may be written together as

$$m_{LR} = \begin{pmatrix} \lambda^m & 0 & 0 \\ 0 & \lambda^n & 0 \\ 0 & 0 & 1 \end{pmatrix} m_f, \quad (4)$$

where m_f corresponds to $(m_\tau \tan \beta)$ in SUSY models for charged lepton mass matrix in case (i), and m_t for up-quark mass matrix in case (ii). The pair of the exponents (m, n) are $(6, 2)$ for charged lepton and $(8, 4)$ for up-quark mass matrices respectively. The value of the Wolfenstein parameter is taken as $\lambda = 0.22$. Using the diagonal form of m_{LR} in Eq.(4) and a suitable choice of the non-diagonal texture of M_{RR} , the following four types of neutrino mass matrices m_{LL} are calculated.

I(A). Nearly degenerate mass matrix with opposite sign mass eigenvalues

The degenerate mass matrix m_{LL} having opposite sign mass eigenvalues is now generated through see-saw formula[8] in Eq.(1) for the choices of m_{LR} in Eq.(4) and

$$M_{RR} = \begin{pmatrix} -2\delta_2 \lambda^{2m} & (\frac{1}{\sqrt{2}} + \delta_1) \lambda^{m+n} & (\frac{1}{\sqrt{2}} + \delta_1) \lambda^m \\ (\frac{1}{\sqrt{2}} + \delta_1) \lambda^{m+n} & (\frac{1}{2} + \delta_1 - \delta_2) \lambda^{2n} & (-\frac{1}{2} + \delta_1 - \delta_2) \lambda^n \\ (\frac{1}{\sqrt{2}} + \delta_1) \lambda^m & (-\frac{1}{2} + \delta_1 - \delta_2) \lambda^n & (\frac{1}{2} + \delta_1 - \delta_2) \end{pmatrix} v_R \quad (5)$$

leading to a simple form,

$$m_{LL} = \begin{pmatrix} -2\delta_1 + 2\delta_2 & \frac{1}{\sqrt{2}} - \delta_1 & \frac{1}{\sqrt{2}} - \delta_1 \\ \frac{1}{\sqrt{2}} - \delta_1 & \frac{1}{2} + \delta_2 & -\frac{1}{2} + \delta_2 \\ \frac{1}{\sqrt{2}} - \delta_1 & -\frac{1}{2} + \delta_2 & \frac{1}{2} + \delta_2 \end{pmatrix} m_0 \quad (6)$$

where m_0 controls the overall magnitude of the masses of the neutrinos whereas δ_1 and δ_2 give the desired splittings for solar and atmospheric data. When $\delta_1 = \delta_2 = 0$, Eq.(6) reduces to the zeroth order mass matrix of the Type I(A) in Table-I, with no splittings[10]. The diagonalisation of m_{LL} in Eq.(6) leads to the following eigenvalues and mixings:

$$m_{\nu_1} = [1 + 2\delta_2 - \delta_1(1 + \sqrt{2})]m_0,$$

$$m_{\nu_2} = [-1 + 2\delta_2 - \delta_1(1 - \sqrt{2})]m_0,$$

$$m_{\nu_3} = m_0,$$

$$\sin^2 2\theta_{12} \approx (1 - \delta_1^2/8), \sin^2 2\theta_{23} = 1, \sin^2 2\theta_{13} = 0$$

For the choice of the values of the parameters $m_0 = 0.4$ eV, $\delta_1 = 0.0061875$,

$\delta_2 = 0.0030625$, Eq.(6) leads to the following numerical predictions

Mixing angles:

$$\sin^2 2\theta_{12} = 0.999, \sin^2 2\theta_{23} \approx 1.0, |V_{e3}| = 6.124 \times 10^{-9},$$

Mass eigenvalues:

$$m_{\nu_i} = (0.396484, -0.396532, 0.4) \text{ eV}, i = 1, 2, 3; \Delta m_{12}^2 = 3.806 \times 10^{-5} eV^2 \text{ and } \Delta m_{23}^2 = 2.76 \times 10^{-3} eV^2.$$

The prediction on solar mixing angle is maximal and larger than the LMA MSW solution. The expression for m_0 in Eq.(6) for case (i) is worked out as $m_0 = m_\tau^2 \tan^2 \beta / v_R$. For input values of $m_0 = 0.4 eV$, $\tan \beta = 40$, $m_\tau = 1.7$ GeV, the see-saw scale is calculated as $v_R \approx 10^{13}$ GeV. This in turn gives the masses of the three right-handed Majorana neutrinos after the diagonalisation of M_{RR} :

$$|M_{RR}^{diag}| = (5.0427 \times 10^{12}, 3.0981 \times 10^8, 1.9613 \times 10^7) \text{ GeV}.$$

Similarly, in case(ii) we have $m_0 = m_t^2 / v_R$ in Eq.(6), and with the input values $m_0 = 0.4 \text{ eV}$, $m_t = 200 \text{ GeV}$, we obtain $v_R = 10^{14} \text{ GeV}$ and the mass eigenvalues of the right-handed Majorana neutrinos: $|M_{RR}^{diag}| = (4.5932 \times 10^{15}, 7.2731 \times 10^6, 5.005 \times 10^{13}) \text{ GeV}$.

I(B). Nearly degenerate mass matrix with the same sign mass eigenvalues

The mass matrix m_{LL} of this type can be realised in the see-saw mechanism(1) using the general texture of m_{LR} in Eq.(4) and

$$M_{RR} = \begin{pmatrix} (1 + 2\delta_1 + 2\delta_2)\lambda^{2m} & \delta_1\lambda^{m+n} & \delta_1\lambda^m \\ \delta_1\lambda^{m+n} & (1 + \delta_2)\lambda^{2n} & \delta_2\lambda^n \\ \delta_1\lambda^m & \delta_2\lambda^n & (1 + \delta_2) \end{pmatrix} v_R \quad (7)$$

leading to the nearly degenerate mass matrix,

$$m_{LL} = \begin{pmatrix} (1 - 2\delta_1 - 2\delta_2) & -\delta_1 & -\delta_1 \\ -\delta_1 & (1 - \delta_2) & -\delta_2 \\ -\delta_1 & -\delta_2 & (1 - \delta_2) \end{pmatrix} m_0 \quad (8)$$

The diagonalisation of m_{LL} in Eq.(8) leads to

$$m_{\nu_1} \simeq (1 - 2\delta_2 - (\sqrt{3} + 1)\delta_1)m_0,$$

$$m_{\nu_2} \simeq (1 - 2\delta_2 + (\sqrt{3} - 1)\delta_1)m_0,$$

$$m_{\nu_3} \simeq m_0,$$

$$\sin^2 2\theta_{12} = \frac{2}{3}, \sin^2 2\theta_{23} = 1, \sin^2 2\theta_{13} = 0.$$

For the choice of the values of the parameters $m_0 = 0.4 \text{ eV}$, $\delta_1 = 3.6 \times 10^{-5}$,

$\delta_2 = 3.9 \times 10^{-3}$, Eq.(8) leads to the following numerical predictions:

Mixing angles:

$$\sin^2 2\theta_{12} = 0.67, \sin^2 2\theta_{23} \approx 1.0, |V_{e3}| = 1.5 \times 10^{-14},$$

Mass eigenvalues:

$$m_{\nu_i} = (0.39684, 0.396892, 0.4) \text{ eV}, i = 1, 2, 3; \Delta m_{12}^2 = 4.13 \times 10^{-5} eV^2 \text{ and } \Delta m_{23}^2 = 2.48 \times 10^{-3} eV^2.$$

The prediction on solar mixing angle is consistent with the LMA MSW solution[3].

For case(i), the expression for m_0 in Eq.(8) is again worked out as $m_0 = m_\tau^2 \tan^2 \beta / v_R$, and for input values of $m_0 = 0.4 eV$, $\tan \beta = 40$, $m_\tau = 1.7 \text{ GeV}$, we obtain $v_R = 1.156 \times 10^{13} \text{ GeV}$ which leads to $|M_R^{diag}| = (1.15 \times 10^{13}, 2.71 \times 10^{10}, 1.498 \times 10^5) \text{ GeV}$. Similarly for case (ii), we have $m_0 = m_t^2 / v_R$ and for input values of $m_0 = 0.4 eV$ and $m_t = 200 \text{ GeV}$, we have $v_R = 1.0 \times 10^{14} \text{ GeV}$ and $|M_{RR}^{diag}| = (1.0039 \times 10^{14}, 5.5089 \times 10^8, 3.035 \times 10^4) \text{ GeV}$.

I(C). Nearly degenerate mass matrix with opposite sign mass eigenvalues

We consider another texture for the nearly degenerate mass matrix m_{LL} with opposite mass eigenvalues[5]. We take m_{LR} given in Eq.(4) and the following right-handed neutrino mass matrix

$$M_{RR} = \begin{pmatrix} (1 + 2\delta_1 + 2\delta_2)\lambda^{2m} & \delta_1 \lambda^{m+n} & \delta_1 \lambda^m \\ \delta_1 \lambda^{m+n} & \delta_2 \lambda^{2n} & (1 + \delta_2)\lambda^n \\ \delta_1 \lambda^m & (1 + \delta_2)\lambda^n & \delta_2 \end{pmatrix} v_R \quad (9)$$

The resulting m_{LL} is calculated as

$$m_{LL} = \begin{pmatrix} (1 - 2\delta_1 - 2\delta_2) & -\delta_1 & -\delta_1 \\ -\delta_1 & -\delta_2 & (1 - \delta_2) \\ -\delta_1 & (1 - \delta_2) & -\delta_2 \end{pmatrix} m_0 \quad (10)$$

where m_0 controls the overall magnitude of the masses of the neutrinos whereas δ_1 and δ_2 give the desired splittings for solar and atmospheric data. When $\delta_1 = \delta_2 = 0$, Eq.(10) reduces to the zeroth order mass matrix of the Type I(C) in Table-I, with no splittings[5,10]. The diagonalisation of m_{LL} in Eq.(10) leads to the following eigenvalues and mixing angles:

$$\begin{aligned} m_{\nu_1} &\simeq (1 - 2\delta_2 - (\sqrt{3} + 1)\delta_1)m_0, \\ m_{\nu_2} &\simeq (1 - 2\delta_2 + (\sqrt{3} - 1)\delta_1)m_0, \\ m_{\nu_3} &\simeq -m_0, \end{aligned}$$

$$\sin^2 2\theta_{12} = \frac{2}{3}, \sin^2 2\theta_{23} = 1, \sin^2 2\theta_{13} = 0.$$

The numerical solution leads to $m_{\nu_i} = (0.39684, 0.396892, -0.4)$ eV, $i = 1, 2, 3$ for the same choices of the input values of $\delta_{1,2}$ and m_0 as in Eq.(8). Further, the predictions on the three mixing angles are the same as in Eq.(8). When $\delta_1 = \delta_2 = 0$, it reduces to the Type I(C) in the Table-I. In case (i), for input values of $m_0 = 0.4$ eV, $\tan\beta = 40$, $m_\tau = 1.7$ GeV, the see-saw scale is calculated as $v_R \approx 10^{13}$ GeV. This in turn gives the masses of the three right-handed Majorana neutrinos after the diagonalisation of M_{RR} : $|M_{RR}^{diag}| = (4.67 \times 10^{11}, 1.296 \times 10^5, 5.06 \times 10^4)$ GeV. Similarly for case (ii), we have $m_0 = m_t^2/v_R$ in Eq.(10), and with the input values $m_0 = 0.4$ eV, $m_t = 200$ GeV, we obtain $v_R = 10^{14}$ GeV and the mass eigenvalues of the right-handed Majorana neutrinos: $|M_{RR}^{diag}| = (1.105 \times 10^{11}, 3.035 \times 10^3, 5.005 \times 10^{11})$ GeV.

II(A). Inverted hierarchical mass matrix with same sign mass eigenvalues

The most general form of the inverted hierarchical mass matrix m_{LL} with the same sign mass eigenvalues, can be calculated with the choice of m_{LR} given in Eq.(4) and M_{RR} of the following form

$$M_{RR} = \begin{pmatrix} 2a\eta(1+2\epsilon)\lambda^{2m} & \eta\epsilon\lambda^{m+n} & \eta\epsilon\lambda^m \\ \eta\epsilon\lambda^{m+n} & a\lambda^{2n} & -(a-\eta)\lambda^n \\ \eta\epsilon\lambda^m & -(a-\eta)\lambda^n & a \end{pmatrix} \frac{v_R}{2a\eta} \quad (11)$$

leading to

$$m_{LL} = \begin{pmatrix} (1-2\epsilon) & -\epsilon & -\epsilon \\ -\epsilon & a & (a-\eta) \\ -\epsilon & (a-\eta) & a \end{pmatrix} m_0 \quad (12)$$

where $a = 0.5$ and m_0 is the overall factor for the masses of the neutrinos. The parameters ϵ and η give the desired splittings for solar and atmospheric data. The diagonalisation of m_{LL} in Eq.(12) leads to the following eigenvalues and mixing angles:

$$m_{\nu_1} \simeq (1 - (\sqrt{3} + 1)\epsilon - \frac{\eta}{2} + \frac{\sqrt{\eta\epsilon}}{6})m_0,$$

$$m_{\nu_2} \simeq (1 + (\sqrt{3} - 1)\epsilon - \frac{\eta}{2} - \frac{\sqrt{\eta\epsilon}}{6})m_0,$$

$$m_{\nu_3} \simeq \eta m_0,$$

and mixing angles:

$$\sin^2 2\theta_{12} = \frac{2}{3}, \sin^2 2\theta_{23} = 1, \sin^2 2\theta_{13} = 0.$$

When $\epsilon = \eta = 0$, Eq.(12) reduces to the zeroth order mass matrix of the

type II(A) in Table-I, with no solar splitting[4,10]. For solution of the LMA MSW solar data and atmospheric neutrino oscillation, we have the choice of the parameters $m_0 = 0.05$ eV, $\epsilon = 0.002$ and $\eta = 0.0001$ leading to the following predictions:

Mixing angles:

$$\sin^2 2\theta_{12} = 0.67, \sin^2 2\theta_{23} \approx 1.0, |V_{e3}| = 3.04 \times 10^{-13},$$

Mass eigenvalues:

$$m_{\nu_i} = (0.05007, 0.04973, 0.000005) \text{ eV}, i = 1, 2, 3; \text{ leading to } \Delta m_{12}^2 = 3.393 \times 10^{-5} \text{ eV}^2 \text{ and } \Delta m_{23}^2 = 2.47 \times 10^{-3} \text{ eV}^2.$$

The expression for m_0 in Eq.(12) for case (i)) is given by $m_0 = m_t^2 \tan^2 \beta / v_R$. For input values of $m_0 = 0.05$ eV, $\tan \beta = 5$, $m_\tau = 1.7$ GeV, we obtain $v_R = 1.445 \times 10^{12}$ GeV which leads to $|M_{RR}^{diag}| = (9.742 \times 10^3, 2.831 \times 10^4, 7.24 \times 10^{15})$ GeV. Again for case(ii) $m_0 = m_t^2 / v_R$ in Eq.(12). Using the input values $m_0 = 0.05 \text{ eV}$, $m_t = 200 \text{ GeV}$, we have $v_R = 8 \times 10^{14}$ GeV and $|M_{RR}^{diag}| = (2.4 \times 10^4, 4 \times 10^{18}, 2.4 \times 10^9)$ GeV where the mass of the heaviest right-handed Majorana neutrino lies above the GUT scale but below the Planck scale[10].

II(B). Inverted hierarchical mass matrix with opposite sign mass eigenvalues

Here the first two neutrino mass eigenvalues are of opposite sign and this inverted hierarchical mass matrix has the following form[11],

$$m_{LL} = \begin{pmatrix} \epsilon & 1 & 1 \\ 1 & \delta_1 & \delta_2 \\ 1 & \delta_2 & \delta_1 \end{pmatrix} m_0, \epsilon, \delta_1, \delta_2 \ll 1 \quad (13)$$

For $\delta_1, \delta_2, \epsilon = 0$, it leads to the type II(B) in Table-I. This structure has been successfully generated within this model in our earlier paper[6], and without much details we outline one case only. As an example, for case (ii) where $m_{LR} = m_{up}$, we have

$$m_{LL} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & \lambda^3 & 0 \\ 1 & 0 & \lambda^3 \end{pmatrix} m_0 \quad (14)$$

for the texture of M_{RR} :

$$M_{RR} = \begin{pmatrix} -\lambda^{22} & \lambda^{15} & \lambda^{11} \\ \lambda^{15} & \lambda^8 & -\lambda^4 \\ \lambda^{11} & -\lambda^4 & 1 \end{pmatrix} v_R \quad (15)$$

The predictions are

$$m_i = (1.4195, -1.4089, 0.0105)m_0, \quad i = 1, 2, 3$$

$$\sin^2 2\theta_{12} = 0.9999, \quad \sin^2 2\theta_{23} \approx 1.0, \quad |V_{e3}| = 1.11 \times 10^{-16},$$

For input $m_0 = 0.05eV$ we get the correct mass eigenvalues and $v_R = 4.0 \times 10^{16} GeV$. This gives

$$|M_{RR}^{diag}| = (3.055 \times 10^{-12}, 2.44 \times 10^{-8}, 1.0)v_R \text{ GeV}.$$

The predictions of the solar mixing angle in all types except in types I(A) and II(B), agree with the LMA MSW solution. The solar mixings predicted from m_{LL} in Eqs.(6) and (14) (Types I(A) and II(B)) are above the upper experimental limit[3] $\sin^2 2\theta_{12} \leq 0.988$ with the best fit value $\sin^2 2\theta_{12} = 0.8163$. Any fine tuning can hardly improve $\sin^2 2\theta_{12}$. One may expect some spectacular changes if the contribution from the diagonalisation of the charged lepton mass matrix having special entries in the 1-2 block, is taken into consideration in the MNS mixing matrix $V_{MNS} = V_{eL}V_{\nu L}^\dagger$. For example, such a charged lepton mass matrix may have the following structure[12,13]

$$m_l = \begin{pmatrix} 0.00256 & -0.01058 & 0 \\ -0.01058 & 0.04596 & 0 \\ 0 & 0 & 1 \end{pmatrix} m_\tau \quad (16)$$

and its diagonalisation leads to the following diagonalisation matrix and eigenvalues:

$$V_{eL} = \begin{pmatrix} -0.97439 & -0.22488 & 0 \\ -0.22488 & 0.97439 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (17)$$

and $m_l^{diag} = (1.182 \times 10^{-4}, 4.840 \times 10^{-2}, 1.0)m_\tau$. With the contribution from charged lepton to the MNS matrix $V_{MNS} = V_{eL}V_{\nu L}^\dagger$, the mixings angles are calculated as

$\sin^2 2\theta_{12} = 0.8517$, $\sin^2 2\theta_{23} = 0.9494$, and $|V_{e3}| = 0.159$ for Type II(B) in Eq.(14), and $\sin^2 2\theta_{12} = 0.8576$, $\sin^2 2\theta_{23} = 0.9495$, and $|V_{e3}| = 0.159$ for Type I(A) in Eq.(6). These results are now consistent with the LMA MSW solution for solar neutrino anomaly and maximal mixing for the atmospheric

neutrino oscillation, along with the CHOOZ constraint $|V_{e3}| \leq 0.16$. The corresponding left-handed neutrino mass matrix in the diagonal basis of the charged lepton mass matrix, is given by relation $m'_{LL} = V_{eL} m_{LL} V_{eL}^\dagger$. For Type II(B) in Eq.(14) we obtain

$$m'_{LL} = \begin{pmatrix} 0.437972 & -0.897698 & -0.9773193 \\ -0.897698 & -0.443296 & -0.230068 \\ -0.9773193 & -0.230068 & -0.005324 \end{pmatrix} m_0 \quad (18)$$

and for Type I(A) in Eq.(6) we have

$$m'_{LL} = \begin{pmatrix} 0.32668 & -0.74163 & -0.57122 \\ -0.74163 & 0.17014 & -0.64185 \\ -0.57122 & -0.64185 & 0.50306 \end{pmatrix} m_0 \quad (19)$$

It is interesting to note that both m'_{LL} in the above Eqs.(18) and (19) have now acquired relatively larger $|m_{ee}|$ consistent with the double beta decay result[1].

A few comments on the stability condition under radiative corrections are in order. The nearly degenerate mass matrices m_{LL} in Eqs.(6),(8) and (10) are found to be unstable under radiative correction in minimal supersymmetric standard model(MSSM). The inverted hierarchical mass matrix with the same mass eigenvalues given in Eq.(12) is also found to be unstable under radiative correction. However, the inverted hierarchical mass matrix given in Eq.(14) with opposite sign mass eigenvalues, is stable under radiative correction [12,14]. The radiative stability of neutrino mass texture for nearly degenerate eigenvalues remains an outstanding problem at the moment[15,16].

3 Summary

In summary, we generate the textures of the nearly degenerate as well as the inverted hierarchical left-handed Majorana neutrino mass matrices from the see-saw formula using the diagonal form of the Dirac mass matrix and non-diagonal form of the right-handed Majorana neutrino mass matrix. This is a continuation of our earlier work where bimaximal mixings are generated from the texture of M_{RR} in case of hierarchical and inverted hierarchical models. The predictions on lepton mixing angles $\sin^2 2\theta_{12} \approx 0.67$, $\sin^2 2\theta_{23} \approx 1.0$ and

$|V_{e3}| \approx 0$ are in excellent agreement with the experimental values in all cases except for types I(A) and II(B). We also get good predictions for Δm_{12}^2 and Δm_{23}^2 which are necessary for the $0\nu\beta\beta$ decays, LMA MSW solar oscillation and atmospheric oscillation data. In all cases the masses of the right-handed Majorana neutrinos are above the weak scale. With the inclusion of the contribution from the diagonalisation of the charged lepton matrix having a special structure in 1-2 block, we improve the prediction on solar mixings to the right order for types I(A) and II(B).

Though the present work is a model independent analysis without using any underlying symmetry, it would serve as a useful guide to building models under the framework of grand unified theories with extended flavour symmetry. In short the present analysis explores the possible origin of the bimaximal neutrino mixings from the texture of right-handed Majorana mass matrices.

References

- [1] H V Klapdor-Kleingrothaus, A Dietz, H L Harney, I V Krivosheina, Mod. Phys. Lett. **A16**(2001)2409.
- [2] Y Fukuda et al, Super-Kamiokande Collaboration, Phys. Rev. Lett. **85**(2000)3999.
- [3] Y Fukuda et al, Super-Kamiokande Collaboration, Phys. Rev. Lett. **86**(2001)5656 and references therein. See also Q R Ahmed et al, SNO Collaboration, Phys. Rev. Lett. **87**(2001) 071301.
- [4] H V Klapdor-Kleingrothaus, U Sarkar, Mod.Phys.Lett.**A16**(2001)2469,**hep-ph/0201224**; N Haba, T.Suzuki, Mod.Phys.Lett.**A17**(2002)865,**hep-ph/0202143**.
- [5] E Ma, Mod.Phys.Lett.**A17**(2002)289, **hep-ph/0201225**
- [6] N Nimai Singh, Mahadev Patgiri,Intl.J.Mod.Phys.**A17**(2002)3629,**hep-ph/0111319**.
- [7] K S Babu, B Dutta, R N Mohapatra, Phys.Lett. **B458**(1999)93, **hep-ph/9904366**; D Falcone, Phys.Lett.**B479**(2000)1, **hep-ph/0111176**.
- [8] M Gell-Mann, P Ramond, R Slansky, in: Supergravity, North-Holland, Amsterdam, 1979; T Yanagida, in : Proc. of the workshop on Unified

- Theory and Baryon number of the Universe, KEK, Japan, 1979; R N Mohapatra, G Senjanovic, Phys. Rev. Lett. **44**(1980)912.
- [9] S F King, N Nimai Singh, Nucl.Phys. **B591**(2000)3; Nucl.Phys. **B596**(2001)81.
 - [10] G Altarelli, F Feruglio, Phys.Rept. **320**(1999)295.
 - [11] An incomplete list: R Barbieri, L Hall, D Smith, A Strumia, N Weiner, JHEP **9812** (1998)17, **hep-ph/9807235**; A S Joshipura, S D Rindani, Eur.Phys. J.**C14**(2000)85; Q Shafi, Z. Tavartkiladze, Phys. Lett. **B482**(2000)145; S F King, N Nimai Singh, Nucl.Phys.**B596**(2001)81; M Patgiri, N Nimai Singh,(to appear in Indian J.Phys.**A**(2003)), **hep-ph/0112123**; K S Babu, R N Mohapatra,Phys.Lett.**B532**(2002)77, **hep-ph/0201176**.
 - [12] M Patgiri, N Nimai Singh,(to appear in Indian J.Phys.**A**(2003)), **hep-ph/0112123**.
 - [13] T Kitabayashi, M.Yasue, Intl.J.Mod.Phys.**A17**(2002)2519, **hep-ph/0112287**
 - [14] P H Chankowski, W Krolikowski, S Pokorski, Phys.Lett. **B473**(2000)109; N Haba, N Okamura, Eur. Phys. J.**C14**(2000)347.
 - [15] M K Parida, C R Das, G Rajsekharan, **hep-ph/0203097**.
 - [16] A S Joshipura, S D Rindani, **hep-ph/0202064**.